

Generalized Conchoids

Boris Odehnal

December 20, 2017

Abstract

We adapt the classical definition of conchoids as known from the Euclidean plane to geometries that can be modeled within quadrics. Based on a construction by means of cross ratios, a generalized conchoid transformation is obtained. Basic properties of the generalized conchoid transformation are worked out. At hand of some prominent examples - line geometry and sphere geometry - the actions of these conchoid transformations are studied. Linear and also non-linear transformations are presented and relations to well-known transformations are disclosed.

Keywords: conchoid transformation, line geometry, sphere geometry, cross ratio, regulus, Dupin cyclide, Laguerre transformation, equiform transformation, inversion.

MSC 2010: 14J26, 51C99, 51N05, 51N15, 51N35, 53A05, 53A17, 74N10, 93B17.

1 Introduction

The well-known construction of conchoids is usually applied to curves in the Euclidean plane \mathbb{R}^2 . Several conchoids of simple and elementary curves in the Euclidean plane are known and have undergone intensive investigations, see for example [4, 6, 10, 11, 16, 22].

The conchoid construction uses a *focus* F and a *directrix* l (with $F \notin l$ in case of a straight line l). Then, a value $d \in \mathbb{R}$ is chosen and the conchoid c_d of l with respect to F is defined as

$$c_d := \{X_d : \overline{X_d X} = d, X_d \in [X, F], \forall X \in l\} \quad (1)$$

where $\overline{X_d X}$ denotes the Euclidean distance




Figure 1: Some conchoids of a straight line l with respect to the focus F .

of the segment $X_d X$ and $[F, X]$ means the line spanned by F and X . A special example is obtained by choosing l as a straight line which yields the one-parameter family of Nikomedes's conchoids. Pascal's limaçon is the conchoid of a circle l with $F \in l$, see [6, 11, 22]. Fig. 1 shows some members from the Nikomedes family. In Euclidean geometry it makes a difference whether the

distance d is equipped with a sign or not. So, the conchoid c_d has either one or two branches depending on whether d is signed or not.

In the following, the mapping $X \mapsto X_d$ shall be called the *conchoid transformation*. It is clearly seen that the conchoid transformation defined via (1) can be applied to arbitrary submanifolds of any metric space. Thus, the conchoid transformation has been applied to surfaces in Euclidean three-space \mathbb{R}^3 in [13, 14, 15, 17, 18] in order to construct new classes of surfaces admitting rational parametrizations, and thus, making them accessible to the algorithms implemented in CAD systems. In [8], a special affine version of a line geometric conchoid transformation was presented. Conchoids on the Euclidean unit sphere were studied from the algebraic and constructive point of view in [12].




Figure 2: The conchoid of a plane with respect to a point is a surface of revolution with Nikomedes's conchoid for its meridian curve.

None of the constructions presented in [13, 14, 15] preserves the type of the geometric object that undergoes the conchoid transformation. The conchoids of planes and spheres become some algebraic surfaces that somehow imitate the features of the

conchoids known from the Euclidean plane, see Fig. 2. Ruled surfaces transform to arbitrary surfaces which, in general, carry maybe only a finite number of straight lines, cf. [13].

In this paper, we adapt the conchoid construction such that it applies to various geometries that can be modeled within quadrics. This is especially the case for the geometry of lines and spheres in three-dimensional spaces which can be modeled within Plücker's and Lie's quadric. So, we are able to find conchoids within certain classes of geometric objects: Lines or spheres can be mapped to lines and spheres. Consequently, ruled surfaces and channel surfaces are transformed to such surfaces. As a by product, rational parametrizations are also preserved.

However, this concept is not restricted to line and sphere geometry, but these are taken as examples in order to show how the generalized conchoid transformation acts. The group of Euclidean motions can be treated in Study's quadric model and also Möbius geometry can be realized on a sphere, further, isotropic geometries and Laguerre geometry also have quadric models and the generalized conchoid transformation can be used there. We shall not discuss these latter four in detail.

Section 2 is dedicated to the generalized conchoid construction and its basic properties. The special cases of line and sphere geometric conchoid transformations are discussed in Sections 3 and 4. In both sections, we treat linear conchoid transformations and special types of quadratic transformations.

2 The generalized conchoid transformation

Let \mathbb{F} be an arbitrary commutative field with $\text{char } \mathbb{F} \neq 2$. Further, let \mathbb{F}^{n+1} be the $(n+1)$ -dimensional vector space on \mathbb{F} and $\mathbb{P}^n(\mathbb{F})$ be the projective space of n dimensions over \mathbb{F}^{n+1} . A quadric $\mathcal{Q} \subset \mathbb{P}^n(\mathbb{F})$ can be defined by prescribing a symmetric bilinear form $\Omega : \mathbb{F}^{n+1} \times \mathbb{F}^{n+1} \rightarrow \mathbb{F}$. With \mathbf{x} we denote the homogeneous coordinate vector of a point $X \in \mathbb{P}^n(\mathbb{F})$ and the equation of the quadric \mathcal{Q} is then $\Omega(\mathbf{x}, \mathbf{x}) = 0$. In the following, lower case bold letters denote the coordinate vectors while capitals denote the points, *i.e.*, the point P has the coordinate vector \mathbf{p} .

Assume that we are given three points P_i ($i \in \{0, 1, 2\}$) in a quadric $\mathcal{Q} \in \mathbb{P}^n(\mathbb{F})$. The line $[P_0, P_1]$ shall not be contained in the quadric and the plane $\pi = [P_0, P_1, P_2]$ shall not be tangent to \mathcal{Q} . Further, $f := \pi \cap \mathcal{Q}$ shall be a regular conic in \mathcal{Q} . Then, there is always a uniquely defined point P_δ that forms the cross ratio $\delta = \text{cr}(P_0, P_1, P_2, P_\delta)$ with P_0, P_1 , and P_2 . Now, we give

Definition 1. *To any triple (P_0, P_1, P_2) of three different points in a quadric $\mathcal{Q} \subset \mathbb{P}^n(\mathbb{F})$ and to any value $\delta \in \mathbb{F} \cup \{\infty\}$ there exists a uniquely defined fourth point P_δ such that $\text{cr}(P_0, P_1, P_2, P_\delta) = \delta$ provided that $[P_0, P_1] \not\subset \mathcal{Q}$, $\pi = [P_0, P_1, P_2]$ is not tangent to \mathcal{Q} , and $\text{char } \mathbb{F} \neq 2$.*

We call P_δ the δ -conchoid transformation of P_2 with respect to the foci P_0 and P_1 .

Later, we will apply the thus generalized conchoid construction to points in ruled

quadrics. Therefore, we do not exclude the case of collinear points P_0, P_1 , and P_2 . Moreover, four points in a quadric do not have to be coplanar in order to assign a cross ratio to them. By means of a stereographic projection to the Gauss plane, we obtain four points that can be identified with four complex numbers and the definition of a cross ratio of these four points is straight forward, see [2].




Figure 3: The generalized conchoid transformation acting on a quadric.

Remark 1. 1. *None of the three points P_0, P_1, P_2 is distinguished. In fact, any two points P_i and P_j ($i \neq j$) out of the three initial points can be considered the foci of the conchoid transformation. Then, P_δ is the conchoid transform of P_k ($k \neq i, j$) with respect to P_i and P_j .*

2. *Changing the roles of the given points does not really change the conchoid transformation. Only the cross ratio δ may turn into one of the six values $\delta, \delta^{-1}, 1 - \delta, (1 - \delta)^{-1}, \delta(\delta - 1)^{-1}, (\delta - 1)\delta^{-1}$.*

3. *In the case $\text{char } \mathbb{F} = 3$, only the cross ratio $\delta = 2$ yields a conchoid trans-*

form $P_\delta \neq P_0, P_1, P_2$. Since any projective line, and thus, any projective conic carries four points in this case, the conchoid transform P_δ of any three (collinear or con-conic) points is the remaining fourth point. Furthermore, 2 is its additive inverse, P_δ makes P_0, P_1, P_2 a harmonic quadruple. Therefore, the conchoid transform P_δ of P_i with respect to P_j and P_k (with $i \neq j, k$ and $j \neq k$) is the harmonic conjugate of P_i with respect to P_j and P_k .

We can give a coordinate representation of the generalized conchoid transformation in $\mathbb{P}^n(\mathbb{F})$ which is helpful for further investigations:

Theorem 1. Let \mathbb{F}^{n+1} be the $(n+1)$ -dimensional vector space on the commutative field \mathbb{F} with $\text{char } \mathbb{F} \neq 2$. Assume that $\Omega : \mathbb{F}^{n+1} \times \mathbb{F}^{n+1} \rightarrow \mathbb{F}^{n+1}$ is a symmetric non-degenerate bilinear form defining a quadric $\mathcal{Q} \subset \mathbb{P}^n(\mathbb{F})$ by $\mathcal{Q} : \Omega(\mathbf{x}, \mathbf{x}) = 0$.

Then, the conchoid transformation $P_2 \mapsto P_\delta$ as explained in Def. 1 can be given in terms of homogeneous point coordinates in $\mathbb{P}^n(\mathbb{F})$ as

$$\mathbf{p}_\delta = \delta(\delta - 1)\Omega_{12}\mathbf{p}_0 + (1 - \delta)\Omega_{02}\mathbf{p}_1 + \delta\Omega_{01}\mathbf{p}_2 \quad (2)$$

where $\delta \in \mathbb{F} \cup \{\infty\}$, \mathbf{p}_i ($i \in \{0, 1, 2\}$) are the homogeneous coordinates of the points P_i , $\Omega_{ij} := \Omega(\mathbf{p}_i, \mathbf{p}_j)$, and neither pair $(\mathbf{p}_i, \mathbf{p}_j)$ is conjugate with regard to \mathcal{Q} (i.e., $\Omega_{ij} \neq 0$).

Proof. We prove Thm. 1 by constructing the coordinate representation given in (2). In the end, we shall arrive at a parametrization of the conic $f = [P_0, P_1, P_2] \cap \mathcal{Q}$ with

$$\text{cr}(P_0, P_1, P_2, P_\delta) = \delta.$$

For that, we observe

$$\begin{aligned} \text{cr}(P_0, P_1, P_2, P_\delta) &= \\ &= \text{cr}(T_{P_0}, [P_0, P_1], [P_0, P_2], [P_0, P_\delta]) = \delta \end{aligned}$$

where T_{P_0} is the tangent of f at P_0 (see Fig. 4), i.e., the intersection of the tangent hyperplane $T_{P_0}\mathcal{Q}$ of \mathcal{Q} at P_0 with the plane $[P_0, P_1, P_2]$. The lines T_{P_0} , $[P_0, P_1]$, $[P_0, P_2]$, $[P_0, P_\delta]$ from the pencil about P_0 establish the stereographic projection $f \rightarrow [P_1, P_2]$ which preserves cross ratios. If further $T_0 := T_{P_0} \cap [P_1, P_2]$ and $P'_\delta := [P_0, P_\delta] \cap [P_1, P_2]$, then $\text{cr}(P_0, P_1, P_2, P_\delta) = \text{cr}(T_0, P_1, P_2, P'_\delta) = \delta$. The point T_0 is found




Figure 4: The stereographic projection $f \rightarrow [P_1, P_2]$ yields a parametrization of the conic f by means of the cross ratio δ .

as the common point of $T_{P_0}\mathcal{Q} : \Omega(\mathbf{p}_0, \mathbf{x}) = 0$ and the line $\mathbf{g}_{12}(\lambda, \mu) = \lambda\mathbf{p}_1 + \mu\mathbf{p}_2$ (with $(\lambda, \mu) \in \mathbb{F}^2 \setminus \{(0, 0)\}$). This yields

$$\mathbf{t}_0 = -\Omega_{02}\mathbf{p}_1 + \Omega_{01}\mathbf{p}_2.$$

Obviously, $\mathbf{t}_0 \neq \mathbf{p}_i$ for $\Omega_{0i} \neq 0$ for $i \in \{1, 2\}$, by assumption. Now, the pairs $(-\Omega_{02}, \Omega_{01})$, $(1, 0)$, $(0, 1)$ are homogeneous coordinates of T_0, P_1, P_2 on the line $[P_1, P_2]$, and thus, we find P'_δ with $\text{cr}(T_0, P_1, P_2, P'_\delta) = \delta$ as

$$\mathbf{p}'_\delta = (1 - \delta)\Omega_{02}\mathbf{p}_1 + \delta\Omega_{01}\mathbf{p}_2.$$

The stereographic projection $[P_1, P_2] \rightarrow \mathcal{Q}$, and thus, also onto f with center P_0 sends P'_δ to P_δ . Since $P_\delta = \mathcal{Q} \cap [P_0, P'_\delta] \setminus \{P_0\}$, we find $\mathbf{p}_\delta = \mathbf{p}'_\delta + \delta(\delta - 1)\Omega_{12}\mathbf{p}_0$ which completes the proof. \square

The coordinate representation

$$\mathbf{p}_\delta : \mathbb{F} \cup \{\infty\} \rightarrow \mathcal{Q}$$

of P_δ given in (2) is a parametrization of the conic $f \subset \mathcal{Q}$. Hence, $\Omega(\mathbf{p}_\delta, \mathbf{p}_\delta) = 0 \forall \delta \in \mathbb{F} \cup \{\infty\}$. We shall call f the *fiber* (conic) of the conchoid transformation.

From the definition of the generalized conchoid transformation and the analytic representation (2), we can easily conclude:

Corollary 1. *The generalized conchoid transformation is involutive if, and only if, the cross ratio equals $\delta = -1$.*

Proof. Geometrically speaking: If P_δ is the conchoid transform of P_2 with $\delta = -1$ and, if transformed with $\delta = -1$ once again, then we obtain P_2 back. \square

Remark 2. *If $\text{char } \mathbb{F} = 3$, then the non-trivial ($\delta \neq 0, 1, \infty$) generalized conchoid transformation is always involutive.*

None of the points \mathbf{p}_i is geometrically distinguished which is expressed in (2) by the fact that the coordinate representation of \mathbf{p}_δ is a trilinear form: It is linear in each \mathbf{p}_i . This gives rise to the following:

Corollary 2. *1. The generalized conchoid transformation on a quadric \mathcal{Q} can be extended to an automorphic collineation κ of \mathcal{Q} .*

2. *The collineation κ has the fixed points P_0 and P_1 corresponding to the two eigenvalues $\delta^2\Omega_{01}$ and Ω_{01} .*

3. *The polar space F of $f := [P_0, P_1]$ with regard to \mathcal{Q} is fixed point wise and corresponds to the eigenvalue $\delta\Omega_{01}$.*

Proof. 1. Consider \mathbf{p}_2 in (2) as the variable point \mathbf{x} . Then, we observe that each summand depends only linearly on \mathbf{x} : The first two coefficients in the linear combination are $\Omega(\mathbf{p}_1, \mathbf{x})$ and $\Omega(\mathbf{p}_0, \mathbf{x})$ and neither is multiplied with \mathbf{x} . Also the last summand depends only linearly on \mathbf{x} , since Ω_{01} is independent of \mathbf{x} . Since $\mathbf{p}_2 = \mathbf{x} \mapsto \mathbf{p}_\delta$ is homogeneous and linear in \mathbf{x} and $\mathbf{p}_\delta \in \mathcal{Q}$ ($\forall \delta \in \mathbb{F} \cup \{\infty\}$), it is an automorphic collineation of \mathcal{Q} .

2. By letting either $\mathbf{p}_2 = \mathbf{p}_0$ or $\mathbf{p}_2 = \mathbf{p}_1$ in (2), we see that the conchoid transformation returns either $\mathbf{p}_\delta = \delta^2\Omega_{01}\mathbf{p}_0$ or $\mathbf{p}_\delta = \Omega_{01}\mathbf{p}_1$.

3. The three-dimensional polar space F of $f = [P_0, P_1]$ with regard to the quadric $\mathcal{Q} : \Omega(\mathbf{x}, \mathbf{x}) = 0$ is given by the homogeneous linear equations

$$F : \Omega(\mathbf{p}_0, \mathbf{x}) = \Omega(\mathbf{p}_1, \mathbf{x}) = 0. \quad (3)$$

With (2), $\mathbf{x} \mapsto \mathbf{x}_\delta$ and reads

$$\begin{aligned} \mathbf{x}_\delta &= \delta(\delta - 1)\Omega_{1\mathbf{x}}\mathbf{p}_0 + \\ &+ (1 - \delta)\Omega_{0\mathbf{x}}\mathbf{p}_1 + \delta\Omega_{01}\mathbf{x} \end{aligned} \quad (4)$$

with $\Omega_{i\mathbf{x}} := \Omega(\mathbf{p}_i, \mathbf{x})$ for $i=0, 1$. Inserting (3) into (4), we infer $\mathbf{x}_\delta = \delta\Omega_{01}\mathbf{x}$ which holds true for all $\mathbf{x} \in \mathbb{F}^{n+1} \setminus \{\mathbf{o}\}$ subject to (3). \square

Remark 3. *The cross ratio itself is a homogeneous coordinate on a projective line or on a conic section (in fact on any rational normal curve). Therefore, we could replace the inhomogeneous parameter δ in (2) by the homogeneous parameter $(d_0, d_1) \in \mathbb{F}^2 \setminus \{(0, 0)\}$ with $\delta = d_1 d_0^{-1}$. We omit this since $d_0 = 0$ causes $\delta = \infty$, $P_\delta = P_0$ which shall be excluded from our considerations.*

3 Line geometric conchoids

3.1 Linear transformations

In this section, we apply the generalized conchoid transformation to the manifold of lines. For that we use the well-known Klein model for the set of lines in a three-dimensional space. Details, exact definitions, properties, and how to compute with Plücker coordinates can be found in [9, 19, 20, 23].

We describe lines L in projective three-space $\mathbb{P}^3(\mathbb{F})$ by Plücker coordinates

$$(\mathbf{l}; \bar{\mathbf{l}}) = (l_1, l_2, l_3; l_4, l_5, l_6) \in \mathbb{F}^6 \setminus \{\mathbf{o}\} \quad (5)$$

which can be made homogeneous (see [9, 19, 20, 23]). Thus, they can be interpreted as coordinates of points in a projective space of five dimensions. From the definition of Plücker coordinates of lines in three-space it is clear that these six-tuples satisfy a quadratic relation:

$$\begin{aligned} M_2^4 : \frac{1}{2}\Omega_L(L, L) &= \langle \mathbf{l}, \bar{\mathbf{l}} \rangle = \\ &= l_1 l_4 + l_2 l_5 + l_3 l_6 = 0. \end{aligned} \quad (6)$$

M_2^4 is a quadric of four dimensions. It is of index two, *i.e.*, the maximum dimension of subspaces contained in M_2^4 equals two. All points on M_2^4 correspond to lines in $\mathbb{P}^3(\mathbb{F})$ and any line can be described by Plücker coordinates satisfying (6). (An affine or a Euclidean specialization is also possible, see [20, 23].)

The polar form $\Omega_L : \mathbb{F}^6 \times \mathbb{F}^6 \rightarrow \mathbb{F}$ of M_2^4 can be used to characterize pairs of lines. Two different lines L and M are coplanar if, and only if, $\Omega_L(L, M) = 0$, *i.e.*, the corresponding points in \mathbb{P}^5 are conjugate with regard to M_2^4 .

Since M_2^4 is a regular hyperquadric in $\mathbb{P}^5(\mathbb{F})$, there is a nine-parameter family of conics in it. We have to distinguish between three types of regular conics in Plücker's quadric: (1) the transversal intersection of a two-dimensional subspace $\mathbb{P}^2(\mathbb{F}) \subset \mathbb{P}^5(\mathbb{F})$; (2a) a conic in a plane $\mathbb{P}_1^2(\mathbb{F}) \subset M_2^4$ (of the first kind); (2b) a conic in a plane $\mathbb{P}_2^2(\mathbb{F}) \subset M_2^4$ (of the second kind). In the following, the conic of type (1) is the most important. The points on a conic of that type correspond to one particular one-parameter family of lines in a ruled quadric, *i.e.*, a regulus. The points on the conics mentioned in the cases (2a) and (2b) correspond to the rulings of a quadratic cone or to the tangents of a (planar) conic, see Fig. 5. Consequently, the




Figure 5: The three types of reguli.

fibers of the line geometric conchoid transformation are reguli and in some cases the rulings of quadratic cones or the tangents of conics. Fig. 6 shows the a typical fiber regulus.




Figure 6: A regular fiber regulus of the line geometric conchoid transformation.

Since the line geometric conchoid transformation maps points on Plücker's quadric to points on Plücker's quadric, it preserves lines, and thus, it maps ruled surfaces to ruled surfaces, and even congruences and complexes of lines (two- and three-dimensional submanifolds of M_2^4) to such.

By assumption, the case of intersecting focal lines L_0 and L_1 is excluded and so we have:

Theorem 2. *Let L_0 and L_1 be two (skew) lines in $\mathbb{P}^3(\mathbb{F})$ and let $\delta \in \mathbb{F} \cup \{\infty\}$ be a certain fixed value. Then, the line geometric conchoid transformation induces a (regular) automorphic collineation κ of M_2^4 that has a fixed line $f \subset \mathbb{P}^5(\mathbb{F})$ and a fixed three-space $F \subset \mathbb{P}^5(\mathbb{F})$ for its axis. F and f are polar with regard to M_2^4 .*

Proof. It is always possible to choose homogeneous coordinates in $\mathbb{P}^3(\mathbb{F})$ such that the Plücker coordinates of the focal lines are

$$L_0 = (1, 0, 0; 0, 0, 0), \quad L_1 = (0, 0, 0; 1, 0, 0),$$

(satisfying $\Omega_L(L_0, L_1) \neq 0$) and $L_2 = (l_1, l_2, l_3; l_4, l_5, l_6)$. We insert into (2) and arrive at

$$L_\delta = (\delta^2 l_1, \delta l_2, \delta l_3; l_4, \delta l_5, \delta l_6). \quad (7)$$

Obviously, the mapping $\kappa : L_2 \mapsto L_\delta$ is a regular collineation with the diagonal matrix $D = \text{diag}(\delta^2, \delta, \delta, 1, \delta, \delta)$ for its coordinate representation. The linear mapping described by D fixes the quadratic form Ω_L , and thus, it is an automorphic collineation of M_2^4 . The eigenvalues are $t_1 = 1$, $t_2 = \delta^2$, and $t_3 = \delta$ with their algebraic multiplicities $\mu(t_1) = \mu(t_2) = 1$ and $\mu(t_3) = 4$. The fixed points corresponding to 1 and δ^2 are L_0 and L_1 , while δ determines the κ -invariant three-space $F : x_1 = x_4 = 0$. The restriction $\kappa|_f$ of κ to the line $f := [L_0, L_1]$ is a hyperbolic projective mapping with two fixed points L_0, L_1 and the coordinate representation $\text{diag}(\delta^2, 1)$. The three-space F is fixed pointwise under κ and is polar to f according to Cor. 2. \square

Remark 4. *In the case of intersecting focal lines L_0 and L_1 , i.e., $\Omega_L(L_0, L_1) = 0$, κ is a projection (singular collineation) onto a one-dimensional subspace of $\mathbb{P}^5(\mathbb{F})$ as can be seen by choosing*

$$L_0 = (1, 0, 0; 0, 0, 0), \quad L_1 = (0, 1, 0; 0, 0, 0),$$

and $L_2 = (l_1, l_2, l_3; l_4, l_5, l_6)$. With (2), we find $L_\delta = (-\delta l_5, l_4, 0; 0, 0, 0)$ provided that $\delta \neq 0$, otherwise L_δ is a single point. Now, $L_2 \mapsto L_\delta$ is a projection onto the subspace $x_2 = x_3 = x_5 = x_6 = 0$. Note that f and F are polar with regard to M_2^4 .

Along a torsal ruling the tangent planes of the ruled surface do not change, see [9, 19].

A ruled surface that consists of torsal rulings only is called *torsal ruled surface* and its parametrization $\mathcal{R}(t)$ satisfies $\Omega_L(\dot{\mathcal{R}}, \dot{\mathcal{R}}) = 0$ besides $\Omega_L(\mathcal{R}, \mathcal{R}) = 0$, both for all $t \in I$.

The term torsal ruled surface covers cylinders, cones, and the surfaces swept by the tangents of a (space) curve (in $\mathbb{P}^3(\mathbb{R})$). Torsal ruled surfaces in Euclidean three-space can be mapped isometrically onto a Euclidean plane, and therefore, these surfaces are called developable. However, torsality is a projective differential geometric property of a ruled surface (see [9, 19, 20, 23]) and we can say:

Corollary 3. *Torsal ruled surfaces are mapped to torsal ruled surfaces under the linear line geometric conchoid transformation.*

Proof. Since torsality of rulings and ruled surfaces is a projective property, it cannot be harmed by the induced automorphic collineation of M_2^4 . According to Thm. 3, the latter is induced by a projective collineation in \mathbb{P}^3 .

We could also prove the corollary by direct calculation. Assume that $\mathcal{R} : I \subset \mathbb{R} \rightarrow M_2^4$ is a curve in M_2^4 (i.e., $\Omega_L(\mathcal{R}, \mathcal{R}) = 0 \forall t \in I$) all of whose rulings are torsal, i.e., $\Omega_L(\dot{\mathcal{R}}, \dot{\mathcal{R}}) = 0$ for all $t \in I$. Then, we compute \mathcal{R}_δ with (2), differentiate once with respect to t , and verify that $\Omega_L(\dot{\mathcal{R}}_\delta, \dot{\mathcal{R}}_\delta) = 0$ on $I \subset \mathbb{R}$. \square

Some examples shall illustrate the action of

the (linear) line geometric conchoid transformation:

Example 1. *The tangents of the curve $(6, 6t, 3t^2, 2t^3) \subset \mathbb{P}^3(\mathbb{R})$ sweep a cubic developable. If we choose $L_0 = (1, 0, 0; 0, 0, 0)$, $L_1 = (0, 0, 0; 1, 0, 0)$ and insert into (2), we obtain the cubic developable built by the tangents of the cubic $(6\delta, 6\delta t, 3t^2, 2t^3)$. If we slice $\mathbb{P}^3(\mathbb{F})$ along $x_0 = 0$ (as usual), we see that the two cubics are related by an affine transformation and so are the cubic developables. Fig. 8 shows the initial cubic developable together with three of its line geometric conchoids. We can see that the*




Figure 8: Some cubic developables being each others line geometric conchoids.

parametrization of the two cubic curves differ only by a multiplication with the matrix $A = \text{diag}(\delta, \delta, 1, 1)$ given in the proof of Thm. 3.

Example 2. *Assume that the focal lines are $L_0 = (0, -1, 2; 0, -2, -1)$ and $L_1 = (1, 1, 3; 3, 0, -1)$. Further, we choose $\delta = 2/5$ and apply the line geometric conchoid*

transformation (2) to the set of rulings given by $L_2(t) = (t^2 + t, t - t^2, 2, -t - t^2, t^2 - t, t^4 + t^2)$ and obtain $L_d(t) = (30t^4 - 25t^2 - 85t - 30, 48t^4 + 151t^2 - 79t - 42, 54t^4 + 63t^2 - 57t - 206, 90t^4 + 205t^2 + 25t - 90, 36t^4 + 2t^2 + 82t - 24, -82t^4 - 79t^2 + 21t + 18)$. A part of this ruled surface is shown in Fig. 9 together with the focal lines and one fiber regulus.




Figure 9: The conchoid transform of a quartic ruled surface with the two focal lines L_0 , L_1 , and a fiber regulus \mathcal{F} .

3.2 Quadratic mappings

The line geometric conchoid transformations discussed in the previous section turned out to be linear mappings, *i.e.*, collineations. From that it is a small step to the definition of a quadratic mapping: Assume that one focal line, say L_1 , is image of L_2 under a fixed projective transformation and leave L_0 fixed. Then, (2) yields a transformation in terms of Plücker coordinates (l_1, \dots, l_6) that is quadratic in the l_i . A special affine version of such a quadratic

line geometric conchoid transformation was studied in [8].

In the following, we shall consider a special Euclidean version of a quadratic line geometric conchoid transformation. For that purpose, we assume that $L_0 = (1, 0, 0; 0, 0, 0)$ is the fixed and constant first focal line. It means no restriction to assume that L_0 coincides with the x -axis of the coordinate system. The second focal line shall be the *absolute polar* of the line $L_2 = (l_1, l_2, l_3; l_4, l_5, l_6)$ (that is to be transformed) with respect to the absolute polarity of Euclidean geometry, *i.e.*, $L_1 = (0, 0, 0; l_1, l_2, l_3)$. With (2) we find the coordinate representation of this particular quadratic line geometric conchoid transformation as

$$q: \begin{pmatrix} l_1 \\ l_2 \\ l_3 \\ l_4 \\ l_5 \\ l_6 \end{pmatrix} \mapsto \begin{pmatrix} \delta(\delta-1)(l_1^2 + l_2^2 + l_3^2) + \delta l_1^2 \\ \delta l_1 l_3 \\ \delta l_2 l_3 \\ (1-\delta)l_1 l_6 + \delta l_3 l_4 \\ (1-\delta)l_2 l_6 + \delta l_3 l_5 \\ l_3 l_6 \end{pmatrix}. \quad (8)$$

The mapping q is degenerate on the field of lines in the ideal plane, *i.e.*, $q(L) = \mathbf{o}$ for all lines $L = (0, 0, 0; u, v, w)$ with $(u, v, w) \in \mathbb{R}^3 \setminus \{\mathbf{o}\}$.

Torsality is, in general, not preserved under quadratic line geometric conchoid transformations. Surprisingly, we can show the following result (for an arbitrarily chosen first focal line L_0) which holds in Euclidean three-space \mathbb{R}^3 :

Theorem 4. *The quadratic line geometric conchoid transformation (8) maps cylinders to cylinders.*

Proof. Let $L_0 = (\mathbf{l}, \bar{\mathbf{l}})$ be the first focal line with constant vectors $\mathbf{l}, \bar{\mathbf{l}} \in \mathbb{R}^3 \setminus \{\mathbf{o}\}$ satisfying $\langle \mathbf{l}, \bar{\mathbf{l}} \rangle = 0$. Further, let

$$L_2 = (\mathbf{v}, \bar{\mathbf{v}}) : I \subset \mathbb{R} \rightarrow M_2^4$$

be the Plücker representation of the cylinder where the constant vector $\mathbf{v} \in \mathbb{R}^3 \setminus \{\mathbf{o}\}$ points into the direction of the cylinder's rulings and $\bar{\mathbf{v}} : I \subset \mathbb{R} \rightarrow \mathbb{R}^3$ is not constant. Naturally, $\langle \mathbf{v}, \bar{\mathbf{v}} \rangle = 0$ for all $t \in I$. Then, the second focal line is given by $L_1 = (\mathbf{o}, \mathbf{v})$ and is obviously constant. Since we are dealing with lines in Euclidean three-space, we may assume that both \mathbf{l} and \mathbf{v} are unit vectors, *i.e.*, $\langle \mathbf{v}, \mathbf{v} \rangle = \langle \mathbf{l}, \mathbf{l} \rangle = 1$.

With $\Omega_{01} = \Omega_{12} = 1$, $\Omega_{02} = \langle \mathbf{l}, \bar{\mathbf{v}} \rangle + \langle \bar{\mathbf{l}}, \mathbf{v} \rangle$, and (2), we find

$$L_\delta = \delta(\delta-1) \begin{pmatrix} \mathbf{l} \\ \bar{\mathbf{l}} \end{pmatrix} + (1-\delta)\Omega_{02} \begin{pmatrix} \mathbf{o} \\ \mathbf{v} \end{pmatrix} + \delta \begin{pmatrix} \mathbf{v} \\ \bar{\mathbf{v}} \end{pmatrix}$$

from which we immediately see that the direction vector

$$\mathbf{l}_\delta = \delta(\delta-1)\mathbf{l} + \delta\mathbf{v} \quad (9)$$

is constant since $\mathbf{l}, \mathbf{v} \in \mathbb{R}^3$ are constant. Therefore, $L_\delta : I \subset \mathbb{R} \rightarrow M_2^4$ parametrizes a cylinder. \square

Remark 5. *Since the argument \mathbf{v} is constant, the quadratic line geometric conchoid transformation turns out to be linear in the case of the cylinder. According to Cor. 3, torsal ruled surfaces are mapped to torsal ruled surfaces.*

It is also possible to verify Thm. 4 via direct computation in terms of coordinates. Then, it is useful to assume that $\mathbf{v} = (0, 0, 1)$ and




Figure 10: The quadratic line geometric conchoid transform of a cylinder.

the cylinder is erected on the cross section $\mathbf{q} = (q_1, q_2, 0) : I \subset \mathbb{R} \rightarrow \mathbb{R}^3$ in the plane $\langle \mathbf{x}, \mathbf{v} \rangle = 0$. Apparently, the cross section of L_δ can be parametrized by

$$\mathbf{q}_\delta = \frac{1-\delta}{\delta l_3} \begin{pmatrix} l_5 \\ -l_4 \\ 0 \end{pmatrix} + \frac{1}{\delta} \begin{pmatrix} q_1 \\ q_2 \\ 0 \end{pmatrix},$$

and thus, the cross section \mathbf{q} of L undergoes an equiform transformation with scaling factor δ^{-1} .

Eq. (9) shows that the direction of the rulings changes. Fig. 10 shows a cylinder of revolution (elliptic cylinder) and its quadratic conchoidal image. The horizontal cross sections of the cylinder are circles and are mapped to circles. The direction of the cylinder's rulings are changed and the image cylinder is again an elliptic cylinder, but with circular horizontal cross sections.

4 Conchoids in sphere geometry

Many of the results from Sec. 3 dealing with lines can be carried over directly to similar results on spheres. This is mainly based on a mapping that goes back to S. LIE and establishes a one-to-one correspondence between lines and spheres. The mapping is called *Lie's line-sphere-mapping* which is a projective collineation $\mathbb{P}^5(\mathbb{C}) \rightarrow \mathbb{P}^5(\mathbb{C})$. Unfortunately, Lie's mapping needs the complex extension of the underlying projective space and mixes up real and complex objects. Therefore, we go a different way and use a coordinatization of the manifold of spheres that was given in [21].

A sphere S in Euclidean three-space \mathbb{R}^3 can always be given by its equation in terms of Cartesian coordinates as

$$S : (s_6 - s_4)(x^2 + y^2 + z^2) - 2s_1x - 2s_2y - 2s_3z + (s_6 + s_4) = 0. \quad (10)$$

For the moment, we assume that $s_6 - s_4 \neq 0$. By completing to full squares in the sphere's equation, we find the center

$$M = \frac{1}{s_6 - s_4}(s_1, s_2, s_3) \quad (11)$$

and the radius R

$$R^2 = \frac{s_1^2 + s_2^2 + s_3^2 + s_4^2 - s_6^2}{(s_6 - s_4)^2}.$$

With the definition

$$R = \frac{s_5}{s_6 - s_4} \quad (12)$$

we find that the six values s_i satisfy the homogeneous quadratic equation

$$L_2^4 : \Omega_S(S, S) := s_1^2 + s_2^2 + s_3^2 + s_4^2 - s_5^2 - s_6^2 = 0. \quad (13)$$

It is clear that the s_i are homogeneous and an interpretation as homogeneous coordinates of points in a projective five-space is nearby. The coordinates $\mathbf{s} = (s_1, \dots, s_6) \in \mathbb{R}^6 \setminus \{\mathbf{o}\}$ are called Lie's sphere coordinates. We shall keep in mind that R can be equipped with a sign which can be used to express an orientation of the sphere S . The quadric L_2^4 spans $\mathbb{P}^5(\mathbb{R})$, is of index one, and therefore, it carries straight lines as maximal subspaces. L_2^4 is called Lie's quadric (cf. [5, 7, 20, 21]) and serves as a point model for the set of spheres in Euclidean three-space.

The polar form $\Omega_S : \mathbb{R}^6 \times \mathbb{R}^6 \rightarrow \mathbb{R}$ describes the polar system of L_2^4 . Assume that S and T are two different spheres (non-proportional Lie coordinates) in oriented contact, *i.e.*, the radii (normal vectors) have equal orientation at the point of contact. Then, S and T are conjugate with regard to L_2^4 , or equivalently, $\Omega_S(S, T) = 0$, and *vice versa*.

The two quadrics M_2^4 and L_2^4 are each others collinear image under Lie's line-sphere-mapping, see [5, 7, 20, 21]. However, the collinear transformation does not map real objects to real ones in general and L_2^4 carries only straight lines, while M_2^4 carries two independent families of planes.

Since L_2^4 carries at most straight lines, there exists only one type of regular conics in L_2^4 . These conics correspond to one-parameter families of spheres enveloping Dupin cyclides. Hence, the fibers of the sphere geometric conchoid transformation are, loosely speaking, Dupin cyclides (cf. Fig. 11). (More precise, but rather lengthy: The fibers of the sphere geometric conchoid




Figure 11: The fibers of the sphere geometric conchoid construction are Dupin cyclides of any type.

transformation are one-parameter families of spheres enveloping Dupin cyclides.)

In analogy to Thm. 2, we can state:

Theorem 5. *Let S_0 and S_1 be two spheres in Euclidean three-space \mathbb{R}^3 (not in oriented contact, i.e., $\Omega_S(S_0, S_1) \neq 0$) and let $\delta \in \mathbb{R} \cup \{\infty\}$ be a certain fixed value. Then, the sphere geometric conchoid transformation induces a (regular) automorphic collineation λ of L_2^4 that has a fixed line f and a fixed three-space L for its axis. F and f are polar with regard to L_2^4 .*

Proof. The proof can be kept short. Without loss of generality, we may assume that the focal spheres are given by

$$S_0 = (0, 0, 0, -1, 1, 0),$$

$$S_1 = (m, 0, 0, \frac{1}{2}(m^2 - R^2 - 1), R, \frac{1}{2}(m^2 - R^2 + 1))$$

are the two focal spheres which are not in oriented contact unless

$$(R - 1 + m)(R - 1 - m) = 0.$$

If now $S_2 = (s_1, \dots, s_6)$ is the sphere to be transformed, then S_δ can be obtained with (2) where Ω is the polar form with the coordinate matrix $\text{diag}(1, 1, 1, 1, -1, -1)$.

In the following, we use the abbreviations

$$\rho_1 = R - m - 1, \quad \rho_2 = R - m + 1.$$

The coordinate matrix of the linear mapping $S_2 \mapsto S_\delta$ has the three different eigenvalues $t_1 = \frac{1}{2}\rho_1\rho_2$, $t_2 = \delta^2 t_1$, and $t_3 = \delta t_1$ with the respective algebraic multiplicities $\mu(t_1) = \mu(t_2) = 1$ and $\mu(t_3) = 4$. Then, it is easily verified that the λ -invariant subspaces show the same behavior as those belonging to κ in the proof of Thm. 2. F and f are polar with regard to L_2^4 according to Cor. 2. \square

Remark 6. *If the focal spheres S_0 and S_1 are in oriented contact, i.e., $\Omega_S(S_0, S_1) = 0$, λ is a projection (singular collineation) onto a one-dimensional subspace of $\mathbb{P}^5(\mathbb{R})$, since then, $(R - 1 + m)(R - 1 - m) = \rho_1\rho_2 = 0$ and the coordinate matrix of $S_2 \mapsto S_\delta$ is of rank 2.*

The above chosen coordinatization of the Euclidean spheres covers more than just Euclidean spheres:

1. (Oriented) Euclidean spheres S are characterized by $s_6 - s_4 \neq 0$ (otherwise the quadratic term in (10) would vanish) and $s_5 \neq 0$, and therefore, $R \neq 0$. Especially, the Euclidean unit sphere S^2 has Lie coordinates $(0, 0, 0, 1, 1, 0)$.
2. (Oriented) planes are characterized by $s_6 - s_4 = 0$. Naturally, the remaining non-vanishing coordinates have to fulfill $s_1^2 + s_2^2 + s_3^2 = s_5^2$. Sometimes, s_5 is set to one and (s_1, s_2, s_3) is then a unit normal vector of the plane

$$\varepsilon : 2s_1x + 2s_2y + 2s_3z - (s_4 + s_6) = 0.$$

3. The hyperplane $s_5 = 0$ meets L_2^4 along the regular three-dimensional quadric $s_1^2 + s_2^2 + s_3^2 + s_4^2 - s_6^2 = 0$ all of whose points correspond to spheres of radius 0. However, spheres of radius 0 can be viewed as points P with coordinates

$$\mathbf{p} = \left(\frac{s_1}{s_6 - s_4}, \frac{s_2}{s_6 - s_4}, \frac{s_3}{s_6 - s_4} \right),$$

but should rather be considered as isotropic cones Γ_P of Euclidean geometry with the equation

$$\Gamma_P : \langle \mathbf{x} - \mathbf{p}, \mathbf{x} - \mathbf{p} \rangle = 0.$$

With $\Gamma_{\mathbf{o}}$ we denote the isotropic cone with the equation $\langle \mathbf{x}, \mathbf{x} \rangle = 0$ emanating from the origin $\mathbf{o} = (0, 0, 0)$ of the coordinate system.

4. Finally, the Lie coordinate vector $(0, 0, 0, 1, 0, 1)$ turns (10) into a false statement, although it describes a point on L_2^4 . It is useful to perform the conformal closure by setting

$$U = (0, 0, 0, 1, 0, 1).$$

Thus, there are four principal types of elements in Lie geometry and any pair out of these four gives rise to a certain sphere geometric conchoid transformation when used as pair of focal spheres. Depending on the nature of the pairs (S_0, S_1) of focal spheres, the sphere geometric conchoid transformations turn out to be well-known transformations from specific subgroups of the huge group of contact transformations. We are able to show the following:

Theorem 6. 1. *The sphere geometric conchoid transformation with $S_0 = U$ and $S_1 = \Gamma_{\mathbf{o}}$ is an equiform transformation, more precisely a similarity with scaling factor δ .*

2. *The sphere geometric conchoid transformation with $S_0 = U$ and $S_1 = S^2$ is a Laguerre transformation.*

3. *The sphere geometric conchoid transformation with $S_0 = \Gamma_{\mathbf{o}}$ and $S_1 = S^2$ is an inversion.*

Proof. 1. It means no restriction to assume that the isotropic cone is centered at the origin of the Cartesian coordinate system. Hence,

$$U = (0, 0, 0, 1, 0, 1),$$

$$\Gamma_{\mathbf{o}} = (0, 0, 0, 1, 0, -1).$$

Inserting $S_0 = U$, $S_1 = \Gamma_{\mathbf{o}}$, and $S_2 = (s_1, \dots, s_6)$ into (2), we find the induced linear mapping

$$\begin{aligned} (s_1, \dots, s_6) &\mapsto (2\delta s_1, 2\delta s_2, 2\delta s_3, \\ &(1 + \delta^2)s_4 + (\delta^2 - 1)s_6, 2\delta s_5, \\ &(\delta^2 - 1)s_4 + (1 + \delta^2)s_6). \end{aligned}$$

Consequently, spheres with center \mathbf{m} and radius ρ are mapped to spheres with center $\delta\mathbf{m}$ and radius $\delta\rho$, while planes are mapped to planes. Further, points are mapped to points. Fig. 12 shows the action of this mapping.




Figure 12: The action of an equiform transformation on one of the one-parameter families of spheres enveloping a torus.

2. In this case, we have $S_0 = U$ (like in the previous case) and

$$S_1 = S^2 = (0, 0, 0, 1, 1, 0).$$

With $S_2 = (s_1, \dots, s_6)$ and (2), we find the induced linear mapping

$$\begin{aligned} (s_1, \dots, s_6) &\mapsto (\delta s_1, \delta s_2, \delta s_3, \\ &\delta(\delta-1)(s_4-s_5) + (\delta-1)s_6 + s_4, \\ &\delta s_5 + (\delta-1)(s_6-s_4), \\ &\delta(\delta-1)(s_4-s_5) + \delta s_6). \end{aligned}$$

Obviously, a sphere with center \mathbf{m} and radius ρ is mapped to a sphere with center $\delta\mathbf{m}$ and radius

$$\frac{\delta s_5 + (\delta-1)(s_6-s_4)}{s_6-s_4} = \delta\rho + \delta - 1.$$

Further, a plane with the equation




Figure 13: A Laguerre transformation is applied to one of the one-parameter families of spheres enveloping a torus.

$s_1x + s_2y + s_3z + s_4 = 0$ with $s_1^2 + s_2^2 + s_3^2 = s_5^2 = 1$ is mapped to the plane $s_1x + s_2y + s_3z + s_4 - 1 + \delta + \delta\rho = 0$ which makes the present sphere geometric conchoid a Laguerre transformation, cf. [3, 5, 7]. Fig. 13 shows how the spheres in a torus change under a Laguerre transformation.

3. Finally, we choose the focal spheres

$$S_0 = \Gamma_{\mathbf{o}} = (0, 0, 0, 1, 0, -1),$$

$$S_1 = S^2 = (0, 0, 0, 1, 1, 0).$$

Thus, (2) yields the linear mapping

$$\begin{aligned} (s_1, \dots, s_6) &\mapsto (\delta s_1, \delta s_2, \delta s_3, \\ &\delta(\delta-1)(s_4-s_5) + (1-\delta)s_6 + s_4, \\ &\delta s_5 + (1-\delta)(s_4+s_6), \\ &\delta(1-\delta)(s_4-s_5) + \delta s_6) \end{aligned}$$

which maps spheres with center \mathbf{m} and radius ρ to spheres with center

$$\frac{\delta\mathbf{m}}{\delta(2\rho\delta - 2\rho + \delta) + (\rho^2 - \langle \mathbf{m}, \mathbf{m} \rangle)(\delta-1)^2}$$




Figure 14: An inversion maps a one-parameter family of spheres enveloping a torus to a one-parameter family of spheres enveloping a Dupin cyclide.

and radius

$$\frac{(\langle \mathbf{m}, \mathbf{m} \rangle - \rho^2)(\delta - 1) - \rho\delta}{(\langle \mathbf{m}, \mathbf{m} \rangle - \rho^2)(\delta - 1)^2 - \delta(2\rho\delta - 2\rho + \delta)}.$$

Obviously, this is an inversion as illustrated in Fig. 14. \square

4.1 Quadratic mappings

Similar to the case of line geometric conchoid transformations, quadratic sphere geometric conchoid transformations can be defined. Therefore, it is only necessary to let the focal sphere S_1 be a linear image of the sphere S_2 to be transformed. A linear image of S_2 means a linear image of the Lie coordinate vector of the sphere S_2 .

We shall have a look at two special types:

Theorem 7. 1. *The quadratic sphere geometric conchoid transformation with the first focal sphere $S_0 = \Gamma_{\mathbf{o}}$ and the second focal sphere S_1 being the polar plane of $(0, 0, 0)$ with respect to*

$S_2 = (s_1, \dots, s_6)$ is a central similarity with center $(0, 0, 0)$ and scaling factor $\frac{1}{2}(1 - 2\delta)\delta^{-2}$, provided that $\delta \neq 0, \frac{1}{2}$

2. *The quadratic sphere geometric conchoid transformation with the first focal sphere $S_0 = S^2 = (0, 0, 0, -1, 1, 0)$ and the second focal sphere S_1 being the radical plane of S_0 and $S_2 = (s_1, \dots, s_6)$ is a cubic transformation, provided that $\delta \neq 0, 1$.*

Proof. 1. The Lie coordinates of the focal spheres are

$$S_0 = (0, 0, 0, -1, 0, 1),$$

$$S_1 = (s_1, s_2, s_3, -\frac{1}{2}(s_4 + s_6), s_5, -\frac{1}{2}(s_4 + s_6))$$

with $s_5^2 = s_1^2 + s_2^2 + s_3^2$. Then, (2) yields

$$\begin{aligned} S_\delta = & ((1 - 2\delta)s_1, (1 - 2\delta)s_2, (1 - 2\delta)s_3, \\ & -\frac{1}{2}(\delta^2(s_4 - s_6) + s_4 + s_6), \\ & (1 - 2\delta)s_5, \\ & \frac{1}{2}(\delta^2(s_6 - s_4) - s_4 - s_6)) \end{aligned}$$

where $s_4 + s_6 \neq 0$ is canceled, since $\Omega_{01} = 2\Omega_{02} = \Omega_{12}(s_6 - s_4)^{-1}$. This can be expressed by means of the original sphere data of S_2 (center $\mathbf{m} = (m_1, m_2, m_3)$ and radius ρ) via

$$\begin{aligned} s_1 &= m_1, s_2 = m_2, s_3 = m_3, \\ s_5 &= \rho, \\ s_4 &= \frac{1}{2}(\langle \mathbf{m}, \mathbf{m} \rangle - \rho^2 - 1), \\ s_6 &= \frac{1}{2}(\langle \mathbf{m}, \mathbf{m} \rangle - \rho^2 + 1) \end{aligned} \quad (14)$$

and gives the center \mathbf{m}_δ and radius ρ_δ of the image spheres:

$$\mathbf{m}_\delta = \frac{1 - 2\delta}{2\delta^2} \mathbf{m}, \quad \rho_\delta = \frac{1 - 2\delta}{2\delta^2} \rho.$$

2. Inserting the Lie coordinates of the two focal spheres

$$S_0 = (0, 0, 0, -1, 1, 0),$$

$$S_1 = (s_1, s_2, s_3, \frac{1}{2}(s_6 - s_4), s_5, \frac{1}{2}(s_6 - s_4)).$$

into (2) yields

$$\begin{aligned} S_\delta = & (s_1((1-2\delta)s_4 - s_6), \\ & s_2((1-2\delta)s_4 - s_6), \\ & s_3((1-2\delta)s_4 - s_6), \\ & \delta((\delta-1)s_4 - s_5)(s_4 - s_6) - (s_4 + s_5)s_6, \\ & s_5((1-2\delta)s_4 - s_6), \\ & \delta((1-\delta)s_4 + s_6)(s_4 - s_6) - (s_4 - s_5)s_5, \end{aligned}$$

which can be reshaped with (14), and finally, (11) and (12) allow us to compute the center and the radius as

$$\mathbf{m}_\delta = \frac{1}{\delta^2} ((1-\delta)(\langle \mathbf{m}, \mathbf{m} \rangle - \rho^2) + \delta) \mathbf{m}$$

$$\rho_\delta = \frac{1}{\delta^2} ((1-\delta)(\langle \mathbf{m}, \mathbf{m} \rangle - \rho^2) + \delta) \rho.$$

Obviously, \mathbf{m}_δ is cubic in the coordinates of \mathbf{m} and ρ_δ is cubic in ρ . \square

5 Conclusion

The projective models of various geometries allow us to generalize the well-known conchoid transformation as long as a quadric model exists and a cross ratio can be defined. So far we haven't dealt with singular quadrics such as Blaschke's cylinder model for isotropic geometries. The subspaces contained in singular quadrics may cause problems for the generalized conchoid construction.

The sphere model of Möbius geometry could also be a playground for generalized conchoid constructions. Nevertheless, cross ratios of four complex numbers can also be defined and give rise to a generalized conchoid construction in the Gauss plane.

Finally, there is one special quadric serving as a point model for the set of Euclidean motions: It is Study's quadric $S_2^6 \subset \mathbb{P}^6(\mathbb{R})$. Conchoid transformations within Study's quadric may generate special Euclidean motions in Euclidean three space.

Acknowledgements

I have to express my sincere thanks to a careful reviewer whose suggestions helped to improve the paper.

References

- [1] A. ALBANO, M. ROGGERO: *Conchoidal transform of two plane curves*. Appl. Algebra Eng. Comm. Comp. **21/4** (2010), 309–328.
- [2] W. BENZ: *Vorlesungen über Geometrie der Algebren*. Springer-Verlag, Berlin-Heidelberg-New York, 1973.
- [3] W. BLASCHKE: *Vorlesungen über Differentialgeometrie III*. Springer-Verlag, Berlin, 1929.
- [4] E. BRIESKORN, H. KNÖRRER: *Plane Algebraic Curves*. Birkhäuser, Basel - Boston - Stuttgart, 2012.

- [5] T.E. CECIL: *Lie sphere geometry*. Springer, New York; 2nd ed. 2008.
- [6] K. FLADT: *Analytische Geometrie spezieller ebener Kurven*. Akademische Verlagsgesellschaft, Frankfurt am Main, 1962.
- [7] O. GIERING: *Vorlesungen über höhere Geometrie*. Vieweg, Braunschweig - Wiesbaden, 1982.
- [8] M. HAMANN, B. ODEHNAL: *Conchoidal ruled surfaces*. Proc. 15th Int. Conf. Geometry & Graphics, Aug. 1–5, 2012, Montreal, article No. 089.
- [9] J. HOSCHEK: *Liniengeometrie*. Bibliographisches Institut, Zürich, 1971.
- [10] J.D. LAWRENCE: *A catalog of special plane curves*. Dover Publications, New York, 1972.
- [11] G. LORIA: *Spezielle algebraische und transzendente ebene Kurven*. B.G. Teubner, Leipzig, 1908.
- [12] B. ODEHNAL: *Conchoids on the Sphere*. KoG **17** (2014), 43–52.
- [13] M. PETERNELL, D. GRUBER, J. SENDRA: *Conchoid surfaces of rational ruled surfaces*. Comp. Aided Geom. Design **28** (2011), 427–435.
- [14] M. PETERNELL, D. GRUBER, J. SENDRA: *Conchoid surfaces of spheres*. Comp. Aided Geom. Design **30** (2013), 35–44.
- [15] M. PETERNELL, L. GOTTHART, J. SENDRA, AND J.R. SENDRA: *Offsets, Conchoids and Pedal Surfaces*. J. Geom. **106** (2015), 321–339.
- [16] J.G. SEMPLE: *Algebraic Curves*. Clarendon Press, Oxford, 1959.
- [17] J.R. SENDRA, J. SENDRA: *An algebraic analysis of conchoids to algebraic curves*. Appl. Algebra Eng. Comm. Comp. **19**/5 (2008), 285–305.
- [18] J. SENDRA, J.R. SENDRA: *Rational parametrization of conchoids to algebraic curves*. Appl. Algebra Eng. Comm. Comp. **21**/4 (2010), 413–428.
- [19] R. SAUER: *Projektive Liniengeometrie*. W. de Gruyter & Co., Berlin und Leipzig, 1937.
- [20] E.A. WEISS: *Einführung in die Liniengeometrie und Kinematik*. B.G. Teubner, Leipzig und Berlin, 1935.
- [21] E.A. WEISS: *Die geschichtliche Entwicklung der Lehre von der Geraden-Kugel-Transformation VII*. Deutsche Math. **3** (1938), 11–35.
- [22] H. WIELEITNER: *Spezielle ebene Kurven*. G.J. Göschen'sche Verlagshandlung, Leipzig, 1908.
- [23] K. ZINDLER: *Liniengeometrie mit Anwendungen I & II*. G.J. Göschen'sche Verlagshandlung, Leipzig, 1902.