

A Spatial Version of the Theorem of the Angle of Circumference

Georg Glaeser¹, Boris Odehnal¹, and Hellmuth Stachel²

¹ University of Applied Arts Vienna, Vienna, Austria
georg.glaeser@uni-ak.ac.at, boris.odehnal@uni-ak.ac.at,

² TU Wien, Vienna, Austria

stachel@dmg.tuwien.ac.at

Abstract. We try a generalization of the theorem of the angle of circumference to a version in three-dimensional Euclidean space and ask for pairs (ε, φ) of planes passing through two (different skew) straight lines $e \ni \varepsilon$ and $f \ni \varphi$ such that the angle α enclosed by ε and φ is constant. It turns out that the set of all such intersection lines is a quartic ruled surface Φ with $e \cup f$ being its double curve. We shall study the surface Φ and its properties together with certain special appearances showing up for special values of some shape parameters such as the slope of e and f (with respect to a fixed plane) or the angle α .

Keywords: ruled surface, angle of circumference, quartic ruled surface, Thaloid, isoptic surface

1 Introduction

The theorem of the angle of circumference states that a straight line segment (bounded by two points E and F) in the Euclidean plane is seen at a constant angle α from any point of a pair of circular arcs passing through E and F . Especially, if the visual angle α is a right angle, the pair of circles becomes one circle with diameter EF , usually referred to as the Thales circle. It would be natural to generalize the theorem of the angle of circumference in Euclidean three-space by asking for all points that see a straight line segment bounded by two points E and F under a constant angle α . The locus of all such points is an algebraic surface of degree four. It is obvious that the latter surface has a rotational symmetry - the axis of the rotation coincides with the straight line $[E, F]$ - and this isoptic surface can be obtained by rotating the pair of circular arcs through E and F , and is therefore, a torus, see Fig. 1. Isoptic curves of spherical conics are also well-known, see [1].

In this contribution, we try a line geometric generalization. We ask for the set of all intersection lines r of planes from two pencils. The lines r can be considered as *one-dimensional eye* seeing a pair of straight lines under a constant angle.

In Section 2, we shall derive the equation of the ruled surface Φ carrying all the lines that see a pair (e, f) of (skew) straight lines under constant angle. From




Fig. 1. A possible generalization of the theorem of the angle of circumference in three-dimensional Euclidean space.

the equation of Φ we can derive some properties of the surface which shall be the contents of Section 3. Finally, in Section 4 we look at special cases of the ruled surface Φ that appear if either the axes e and f reach a special relative position or if the angle α attains special values or if even both is the case.

2 Equation of the ruled surface

It is favorable to represent points in Euclidean three-space \mathbb{R}^3 by Cartesian coordinates (x, y, z) . It means no restriction to assume that the axes e and f of the two pencils of planes are given by

$$e = \begin{pmatrix} d \\ 0 \\ 0 \end{pmatrix} + t \begin{pmatrix} 0 \\ 1 \\ k \end{pmatrix}, f = \begin{pmatrix} -d \\ 0 \\ 0 \end{pmatrix} + u \begin{pmatrix} 0 \\ 1 \\ -k \end{pmatrix} \quad (1)$$

parametrized by real parameters $t, u \in \mathbb{R}$. Here and in the following, $\overline{ef} = 2d \in \mathbb{R}$ is the distance between the straight lines e, f and $k \in \mathbb{R}$ is their slope with respect to the plane $z = 0$, see Fig. 2. Since $\mathbf{g} = (0, 1, k)$ and $\mathbf{h} = (0, 1, -k)$ are the directions of the lines e and f , the normals of the planes $\varepsilon \ni e$ and $\varphi \ni f$ are linear combinations of $\mathbf{g}_1 = (0, -k, 1)$, $\mathbf{g}_2 = (1, 0, 0)$ or $\mathbf{h}_1 = (0, k, 1)$, $\mathbf{h}_2 = \mathbf{g}_2$, respectively. With $\lambda, \mu \in \mathbb{R}$ we let

$$\mathbf{n}_\varepsilon = \mathbf{g}_1 + \lambda \mathbf{g}_2, \quad \mathbf{n}_\varphi = \mathbf{h}_1 + \mu \mathbf{h}_2. \quad (2)$$

Now, we can write down the condition $\sphericalangle(\varepsilon, \varphi) = \sphericalangle(\mathbf{n}_\varepsilon, \mathbf{n}_\varphi) = \alpha$ by evaluating

$$\langle \mathbf{n}_\varepsilon, \mathbf{n}_\varphi \rangle^2 = A^2 \langle \mathbf{n}_\varepsilon, \mathbf{n}_\varepsilon \rangle \langle \mathbf{n}_\varphi, \mathbf{n}_\varphi \rangle$$




Fig. 2. Choice of a Cartesian coordinate system and the meaning of d and k .

where $\langle \mathbf{u}, \mathbf{v} \rangle$ denotes the canonical scalar product of two vectors $\mathbf{u}, \mathbf{v} \in \mathbb{R}^3$ and $A := \cos \alpha$. This gives

$$(1 - k^2 + \lambda\mu)^2 = A^2(1 + k^2 + \lambda^2)(1 + k^2 + \mu^2). \quad (3)$$

The planes from either pencil have normal vectors given in (2), and thus, they have the equations

$$\begin{aligned} \varepsilon : \lambda x - ky + z &= d\lambda, \\ \varphi : \mu x + ky + z &= -d\mu. \end{aligned} \quad (4)$$

So far, both λ and μ can vary freely in \mathbb{R} , and thus, the planes ε and φ intersect in the lines of a hyperbolic linear line congruence with axes e and f , parametrized by

$$\mathbf{r}(t, \lambda, \mu) = \frac{d}{k} \begin{pmatrix} k \\ -\mu \\ -\mu k \end{pmatrix} + t \begin{pmatrix} -2k \\ \mu - \lambda \\ k(\lambda + \mu) \end{pmatrix} \quad (5)$$

where $t \in \mathbb{R}$ is the parameter on the lines in the congruence. The ruled surface Φ we are aiming at is precisely that subset of the congruence (5) where λ and μ are subject to (3).

The equation of the ruled surface Φ in terms of Cartesian coordinates is obtained from the parametrization (5) by eliminating all parameters t, λ, μ : Assume $\mathbf{r} = (r_x, r_y, r_z)$. Then, we eliminate t from $x - r_x, y - r_y$, and $z - r_z$ by computing the resultants

$$\begin{aligned} r_1 &:= \text{res}(x - r_x, z - r_z, t), \\ r_2 &:= \text{res}(y - r_y, z - r_z, t). \end{aligned}$$

In the next step, we eliminate λ from both, r_1 and r_2 using (3) which results in two further polynomials $r'_1 \in \mathbb{R}[x, z, \mu]$ and $r'_2 \in \mathbb{R}[y, z, \mu]$. It would make no difference if we eliminate μ first. Finally, the resultant of r'_1 and r'_2 with respect

to μ contains a non-trivial factor which is the equation of Φ . (The trivial factors of the latter resultant are detected by substituting (5) and verifying that they do not vanish.)

So, we obtain the following equation of Φ :

$$\begin{aligned} & \sigma_1\sigma_2(x^2 - d^2)^2 - B^2(z^2 - k^2y^2)^2 + \\ & + 2\sigma_3(d^2z^2 + k^2x^2y^2) + 2\sigma_4(d^2k^2y^2 + x^2z^2) + \\ & - 8A^2dk(1 + k^2)xyz = 0 \end{aligned} \quad (6)$$

with the abbreviations

$$\sigma_{1,2} := A k^2 \pm k^2 + A \mp 1, \quad \sigma_{3,4} := A^2k^2 \mp k^2 + A^2 \pm 1,$$

and $B^2 = 1 - A^2$ (or $B = \sin \alpha$). Summarizing, we can state:

Theorem 1. *The isoptic ruled surface Φ as the set of intersection lines r of planes ε, φ from two pencils with axes e, f and $\sphericalangle(\varepsilon, \varphi) = \alpha = \text{const.}$ is the algebraic ruled surface with the equation (6) and is, in general, of degree four.*

Fig. 3 shows an example of a ruled surface Φ together with the axes e and f of the pencils of planes.

In the case $A = 0$ which is equivalent to $\alpha = \frac{\pi}{2}$, there exists a generation of Φ by means of a projective mapping κ from the pencil of planes about e to the pencil of planes about f . The projectivity κ assigns to each plane ε (through e) precisely one plane φ (through f) such that $\varepsilon \perp \varphi$. Thus, the lines $\varepsilon \cap \kappa(\varepsilon)$ form a regulus, *i.e.*, one family of straight lines on a (regular) ruled quadric. Inserting $A = 0$ into (6) returns the equation of the regular ruled quadric which is in any case a hyperboloid (with multiplicity two):

$$((1 - k^2)x^2 - k^2y^2 + z^2 + d^2(k^2 - 1))^2 = 0 \quad (7)$$

an example of which is shown in Fig. 4.

3 Properties of Φ

From the construction of Φ it is clear that the lines e and f are part of the surface. Moreover, the union of these lines is the double curve of Φ . Hence, Φ is of Sturm type 1, cf. [2]. Since Φ is of Sturm type 1, it is elliptic.

Each plane ε in the pencil about e intersects Φ along e with multiplicity 2. Since each such plane ε contains at least one generator, the remaining part of $\varepsilon \cap \Phi$ has to be a straight line s too. The line s is a further generator of Φ . A similar statement can be made about the planes φ through f .

The tangent planes of Φ meet Φ along (planar) cubic curves which are either rational or elliptic, see Fig. 5. The quartic ruled surface Φ carries no regular conic: Any plane ε through a pair of intersecting generators g_1, g_2 shares one




Fig. 3. The quartic *isoptic* ruled surface of a pair of skew straight lines e and f .

of the axes, say e , with Φ . Therefore, the remaining part of $\Phi \cap \varepsilon \setminus \{e, g_1, g_2\}$ has to be a straight line l and $l \cup e$ is a singular conic, cf. [2]. If we perform the projective closure of the Euclidean three-space, then we can look Φ 's intersection Φ_∞ with the ideal plane, see Fig. 6. The ideal points E_∞ and F_∞ of the straight lines e and f are the only double points of the elliptic quartic Φ_∞ . An equation of Φ_∞ can be obtained from (6) by removing all terms of degree three and less:

$$\Phi_\infty : (\sigma_1\sigma_2x^2 + 2k^2\sigma_3y^2 + 2\sigma_4z^2)x^2 - B^2(k^2y^2 - z^2)^2 = 0. \quad (8)$$

Then, we interpret $x : y : z$ as homogeneous coordinates of points in the plane at infinity and note that in Φ_∞ 's equation d does not show up.

4 Special cases

We can expect exceptional appearances of the quartic ruled surface Φ if we choose special values for A , d , or k . Although all these values are originally assumed to be real and especially $|A| \leq 1$, we need not restrict ourselves to real values. In the following, we shall discuss some of these special choices that lead to sometimes unexpected surfaces Φ which, eventually, are then no longer ruled surfaces with real rulings.




Fig. 4. A one-sheeted hyperboloid appears if $A = 0$, $d, k \in \mathbb{R}^*$, cf. (7).

4.1 One-sheeted hyperboloids

Intersecting lines \overline{ef} e and f . In the very beginning, we made the natural assumption $2d = \overline{ef} \neq 0$, *i.e.*, the lines e and f are skew. If we allow $d = 0$, then (6) simplifies to (8) which comes as no surprise, since (8) is independent of d . Therefore, (8) can also be viewed as the equation of a quartic cone Γ emanating from $(0, 0, 0)$. Obviously, e and f are generators with multiplicity two. The cone Γ is the asymptotic cone of Φ an example of which is displayed in Fig. 7.

Further, if $k = 0$ (together with $d = 0$ this actually means $e = f$), then Γ degenerates and becomes the pair of isotropic planes $x^2 + z^2 = 0$ with multiplicity two.

If we allow $A = 0$, the cone Γ becomes the quadratic cone

$$(k^2 - 1)x^2 + k^2y^2 - z^2 = 0 \quad (9)$$

with multiplicity two. The quadratic cone (9) is a *normal cone* (cf. [1, p. 463–467]) and it is the asymptotic cone of the double hyperboloid (7) being the special form of Φ if $A = 0$.

An interesting case occurs if $k = \pm i$ (besides $d = 0$), *i.e.*, the axes e and f of the pencils of planes are isotropic lines. In this case, Φ splits into two singular quadrics:

$$\begin{aligned} 2x^2 + (1 - A)y^2 + (1 - A)z^2 &= 0, \\ 2x^2 + (1 + A)y^2 + (1 + A)z^2 &= 0. \end{aligned} \quad (10)$$




Fig. 5. The quartic ruled surface Φ carries a two-parameter family of cubic curves which come as the intersection of Φ with its tangent planes. Here: $\tau \cap \Phi = r \cup c$ where $r \subset \tau$ is a ruling, τ is a tangent plane through r , and c is the cubic curve.




Fig. 6. The intersection of Φ with the plane at infinity is an elliptic quartic curve Φ_∞ with the two double points E_∞ and F_∞ which are the ideal points of Φ 's double curve $e \cup f$.




Fig. 7. Intersecting lines e and f produce a quartic cone which is also the asymptotic cone of the generic (non-degenerate) quartic ruled surface Φ .

One of these becomes a plane with multiplicity two if either $A = +1$ or $A = -1$ while the other one becomes an isotropic cone $x^2 + y^2 + z^2 = 0$.

The case $|A| < 1$ turns both of the quadrics into cones without any real points besides the common vertex $(0, 0, 0)$.

$|A| > 1$ corresponds to purely imaginary angles

$$\alpha \equiv i \cdot \ln(A + \sqrt{A^2 - 1}) \pmod{2\pi}.$$

Nevertheless, inserting $|A| > 1$ into (10) makes either the first or the second quadric a cone with real points while the other still has only one real point, namely the vertex $(0, 0, 0)$.

Parallel lines e and f . The case of parallel axes e and f is clearly an extrusion of the planar figure of the theorem of the angle of circumference. Thus, the ruled surface Φ (6) will split into two cylinders Δ_1 and Δ_2 of revolution erected on those circular arcs in the $[x, y]$ -plane which are the locus of all points seeing the line segment EF (with $E, F = (\pm d, 0, 0)$) under constant angle α (cf. Fig. 8).

From (6) we find the equation of the degenerate quartic by replacing k with $1/K$ and subsequently setting $K = 0$. (Otherwise, we would have to set $k = \infty$.) This results in the expected pair of cylinders of revolution

$$\Delta_{1,2} : x^2 + y^2 \pm \frac{2dA}{\sqrt{1 - A^2}}y - d^2 = 0.$$




Fig. 8. A pair of cylinders of revolution as the isoptic ruled surface of parallel lines e and f .

In order to find real surfaces Δ_i , the values for A are restricted to $|A| \leq 0$. The Thaloid $x^2 + y^2 = d^2$ (cylinder of revolution) through e and f cannot be obtained directly from the cylinders' equations, since then $A = 1$.

Other quadrics. The axes e and f of the pencils of planes can be chosen as isotropic lines. Therefore, we let $k = i$. (The choice $k = -i$ produces the same result.) Again, we find that (6) degenerates and splits into quadratic polynomials:

$$\begin{aligned} Q_1 : 2x^2 + (1 - A)y^2 + (1 - A)z^2 &= 2d^2, \\ Q_2 : 2x^2 + (1 + A)y^2 + (1 + A)z^2 &= 2d^2. \end{aligned} \tag{11}$$

The case $d = 0$ was discussed earlier, so we have $d \neq 0$ in the following. Independent of the choice of A and regardless of the regularity, both quadrics Q_1 and Q_2 have the x -axes for their common axis of revolution.

In the very special case $A = \pm 1$, the pair of quadrics (11) contains precisely the singular quadric $x^2 - d^2 = (x - d)(x + d) = 0$ (a pair of (real) parallel planes) and the Euclidean sphere $x^2 + y^2 + z^2 = d^2$ with radius d centered at $(0, 0, 0)$ touching the planes at $(\pm d, 0, 0)$.

If $|A| > 1$, the pair (Q_1, Q_2) of quadrics consists of a two-sheeted hyperboloid of revolution and an ellipsoid of revolution.

Finally, we obtain two ellipsoids of revolution if $|A| < 1$.

The following table summarizes the special and degenerate cases of Φ depending on special choices of A, k, d .

$A = 0$		
$k = 0$	$k = i$	$k = \infty$
$(d^2 - x^2 - z^2)^2 = 0$ right cylinder, $\mu = 2$	$(2d^2 - 2x^2 - y^2 - z^2)^2 = 0$ ellipsoid, $\mu = 2$	$(d^2 - x^2 - y^2)^2 = 0$ right cylinder, $\mu = 2$
$d = 0$	$d = 0$	$d = 0$
$(x^2 + z^2)^2 = 0$ compl. conj. planes, $\mu = 2$	$(2x^2 + y^2 + z^2)^2 = 0$ cone, no real point $\neq (0, 0, 0)$, $\mu = 2$	$(x^2 + y^2)^2 = 0$ compl. conj. planes, $\mu = 2$
$d = i$	$d = i$	$d = i$
$(1 + x^2 + z^2)^2 = 0$ right cylinder, no real points, $\mu = 2$	$(2x^2 + y^2 + z^2 + 2)^2 = 0$ ellipsoid, no real point, $\mu = 2$	$(1 + x^2 + y^2)^2 = 0$ right cylinder, no real point, $\mu = 2$
$A = 1$		
$k = 0$	$k = i$	$k = \infty$
$z^2 = 0$ plane, $\mu = 2$	$(x^2 - d^2)(d^2 - x^2 - y^2 - z^2) = 0$ sphere \cup tangent planes	$y^2 = 0$ plane, $\mu = 2$
$d = 0$	$d = 0$	$d = 0$
empty set	$x^2(x^2 + y^2 + z^2) = 0$ real double plane \cup \cup isotropic cone	empty set
$d = i$	$d = i$	$d = i$
$z^2 = 0$ plane, $\mu = 2$	$(x^2 + 1)(1 + x^2 + y^2 + z^2) = 0$ sphere, no real point \cup \cup compl. tang. planes	$y^2 = 0$ plane, $\mu = 2$

Table 1. Special shapes of Φ caused by $A = 0, 1$, $\alpha = 0, \frac{\pi}{2}$, $d = 0$, $k = 0, \infty, i$; the integer μ denotes the multiplicities of the components.

References

1. G. GLAESER, H. STACHEL, B. ODEHNAL: *The Universe of Conics*. From the ancient Greeks to 21st century developments. Springer-Spektrum, Springer-Verlag, Heidelberg, 2016.
2. E. MÜLLER: *Vorlesungen über Darstellende Geometrie*. III. Band: Konstruktive Behandlung der Regelflächen. Deuticke, Leipzig und Wien, 1931.
3. H. WIELEITNER: *Spezielle Ebene Kurve*. Sammlung Schubert LVI, G.J. Göschen'sche Verlagshandlung, Leipzig, 1908.